

ON A CLASSIFICATION OF WARPED PRODUCT SPACES WITH GRADIENT RICCI SOLITONS

SANG DEOK LEE, BYUNG HAK KIM*, AND JIN HYUK CHOI

ABSTRACT. In this paper, we study Ricci solitons, gradient Ricci solitons in the warped product spaces and gradient Yamabe solitons in the Riemannian product spaces. We obtain the necessary and sufficient conditions for the Riemannian product spaces to be Ricci solitons. Moreover we classify the warped product space which admit gradient Ricci solitons under some conditions of the potential function.

1. Introduction

A Riemannian metric g on a complete Riemannian manifold M is called a Ricci soliton if there exists a smooth vector field X such that the Ricci curvature tensor satisfies

$$(1) \quad Ric + \frac{1}{2}\mathfrak{L}_X g = \rho g$$

for some constant ρ , where \mathfrak{L}_X is the Lie derivative with respect to X [2, 3, 5, 6, 9]. It is said that (M, g) or M is a Ricci soliton if the metric g on M is a Ricci soliton. Evidently we see that an Einstein metric becomes a Ricci soliton, but the converse is not true. The Ricci

Received November 9, 2016. Revised December 6, 2016. Accepted December 7, 2016.

2010 Mathematics Subject Classification: 53C25, 53B21.

Key words and phrases: Ricci curvature; warped product space; Ricci soliton.

This research was supported by a grant from Kyung Hee University in 2013(KHU-20130566).

* Corresponding author.

© The Kangwon-Kyungki Mathematical Society, 2016.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution and reproduction in any medium, provided the original work is properly cited.

soliton is called shrinking if $\rho > 0$, steady if $\rho = 0$ and expanding if $\rho < 0$. The metric of a Ricci soliton is useful in not only physics but also mathematics, and often referred as quasi-Einstein [?]. If $X = \nabla h$ for some function h on M , then M is called a gradient Ricci soliton. The function h above is called the potential function. In this case, the equation (1) can be rewritten as

$$(2) \quad Ric + Hess h = \rho g.$$

It is well known that when $\rho \leq 0$ all compact solitons are necessarily Einstein [6], and a Ricci soliton on a compact manifold has a constant curvature in 2-dimension [?] as well as in 3-dimension [8]. Moreover a Ricci soliton on a compact manifold is a gradient Ricci soliton [9], and a compact shrinking soliton is always gradient [13].

In [11], we studied the gradient Ricci soliton in the warped product space and proved that the base space of the warped product space with a gradient Ricci soliton to be a gradient Ricci soliton or an Einstein space is determined by the second derivative of the warping function. As far as we know, there are no results for the warped products of two Riemannian manifolds with general dimension and gradient Ricci soliton related to potential function. In this sense, the considerations of the warped products with gradient Ricci soliton satisfying some conditions related to potential function are meaningful. Related these problems, we get some useful results and classification theorem (see Theorem 3.3).

A Riemannian metric g on a Riemannian manifold M is called a Yamabe soliton if there exist a smooth vector field X and a constant ρ such that

$$(3) \quad (r - \rho)g = \frac{1}{2} \mathcal{L}_X g,$$

where r is the scalar curvature of M . In particular, if $X = \nabla h$ for some smooth function h , we call it the gradient Yamabe soliton [?]. The function h above is called the potential function. In this case, the equation (3) can be rewritten as

$$(4) \quad (r - \rho)g = \nabla^2 h.$$

In Chapter 4, we studied gradient Yamabe soliton in $M = R^1 \times B$ for a Riemannian manifold B , and we get the necessary and sufficient condition for M admits a gradient Yamabe soliton.

2. Ricci solitons in the Riemannian product manifolds

Let (B, g) be an m -dimensional Riemannian manifold with a metric g and let $M = R \times B$ be the product Riemannian manifold with the metric $\tilde{g} = \begin{pmatrix} 1 & 0 \\ 0 & g \end{pmatrix}$. Then the Ricci curvature tensors \tilde{S} and S of M and B , respectively are given by $S_{ab} = S_{ab}$ and the others are zero, where the range of indices a, b, c, \dots is $\{2, 3, \dots, m + 1\}$.

Suppose that B is a Ricci soliton. Then there exists a smooth vector field $U = (\xi_2, \dots, \xi_{m+1})$ on B such that

$$S_{ab} = \rho g_{ab} - \frac{1}{2}(\nabla_a \xi_b + \nabla_b \xi_a),$$

for some constant ρ .

If we take $\tilde{\rho} = \rho$ and $\tilde{U} = (\tilde{\xi}_1, \tilde{\xi}_2, \dots, \tilde{\xi}_{m+1}) = (\rho t, \xi_2, \dots, \xi_{m+1})$, then we obtain

$$\begin{aligned} \tilde{S}_{ab} &= S_{ab} = \rho g_{ab} - \frac{1}{2}(\nabla_a \xi_b + \nabla_b \xi_a) = \tilde{\rho} \tilde{g}_{ab} - \frac{1}{2}(\tilde{\nabla}_a \tilde{\xi}_b + \tilde{\nabla}_b \tilde{\xi}_a), \\ (5) \quad \tilde{S}_{a1} &= 0 = -\frac{1}{2}(\tilde{\nabla}_a \tilde{\xi}_1 + \tilde{\nabla}_1 \tilde{\xi}_a), \\ \tilde{S}_{11} &= 0 = \tilde{\rho} - \frac{1}{2}(\tilde{\nabla}_1 \tilde{\xi}_1 + \tilde{\nabla}_1 \tilde{\xi}_1). \end{aligned}$$

That is, M becomes a Ricci soliton.

Conversely, if we suppose that M is a Ricci soliton, then there exists a smooth vector field $\tilde{V} = (\tilde{\xi}_1, \tilde{\xi}_2, \dots, \tilde{\xi}_{m+1})$ on M such that

$$\tilde{S}_{ij} = \tilde{\rho} \tilde{g}_{ij} - \frac{1}{2}(\tilde{\nabla}_i \tilde{\xi}_j + \tilde{\nabla}_j \tilde{\xi}_i)$$

for some constant function $\tilde{\rho}$ on M , where the range of indices i, j, k, \dots is $\{1, 2, 3, \dots, m + 1\}$. So we have

$$\begin{aligned} \tilde{S}_{ab} &= \tilde{\rho} \tilde{g}_{ab} - \frac{1}{2}(\tilde{\nabla}_a \tilde{\xi}_b + \tilde{\nabla}_b \tilde{\xi}_a) = S_{ab}, \\ (6) \quad \tilde{S}_{a1} &= -\frac{1}{2}(\tilde{\nabla}_a \tilde{\xi}_1 + \tilde{\nabla}_1 \tilde{\xi}_a) = -\frac{1}{2}(\partial_a \tilde{\xi}_1 + \partial_1 \tilde{\xi}_a) = 0, \\ \tilde{S}_{11} &= \tilde{\rho} - \frac{1}{2}(\tilde{\nabla}_1 \tilde{\xi}_1 + \tilde{\nabla}_1 \tilde{\xi}_1) = \tilde{\rho} - \partial_1 \tilde{\xi}_1 = 0. \end{aligned}$$

From the third equation of (6), we can put

$$(7) \quad \tilde{\xi}_1 = \tilde{\rho} t + h(x_2, \dots, x_{m+1})$$

for $t \in R$ and some function h on B . Using the second equation of (6) and (7), we see that $\tilde{\xi}_a$ is of the form

$$(8) \quad \tilde{\xi}_a = -h_a t + k^a,$$

where $h_a = \partial_a h$ and $k^a = k^a(x_2, \dots, x_{m+1})$ is a function on B corresponding to $\tilde{\xi}_a$. Hence, the vector field \tilde{V} on M is given by

$$\tilde{V} = (\tilde{\rho}t + h, -h_2 t + k^2, \dots, -h_{m+1} t + k^{m+1}).$$

From the first equation of (6) and the fact that $S_{ab} - \rho g_{ab}$ depends only on B , we find that the equation $S_{ab} - \rho g_{ab} = -\frac{1}{2}(\nabla_a \xi_b + \nabla_b \xi_a)$ holds for $\rho = \tilde{\rho}$ and $\xi_a = k^a$. If we take the vector field $\tilde{V} = (\xi_2, \dots, \xi_{m+1})$ on B , then we see that B is a Ricci soliton. Thus, we have

THEOREM 2.1. *$M = R \times B$ is a Ricci soliton if and only if B is a Ricci soliton.*

Next, consider the case of $M = R^2 \times B$ be the product Riemannian manifold with the Riemannian metric $\tilde{g} = \begin{pmatrix} \delta_{uv} & 0 \\ 0 & g_{ab} \end{pmatrix}$, where the range of indices u, v is $\{1, 2\}$ and the range of indices a, b, c, \dots is $\{3, 4, \dots, m+2\}$. Then, the Ricci curvature tensors \tilde{S} and S of M and B , respectively, are given by $\tilde{S}_{ab} = S_{ab}$ and the others are zero.

Suppose that B is a Ricci soliton. Then there exists a smooth vector field $U = (\xi_3, \dots, \xi_{m+2})$ on B such that $S_{ab} - \rho g_{ab} = -\frac{1}{2}(\nabla_a \xi_b + \nabla_b \xi_a)$ for some constant ρ . Take $\tilde{\rho} = \rho$ and $\tilde{U} = (\tilde{\xi}_1, \tilde{\xi}_2, \dots, \tilde{\xi}_{m+2}) = (\rho t_1, \rho t_2, \xi_3, \dots, \xi_{m+2})$, then we obtain

$$(9) \quad \begin{aligned} \tilde{S}_{ab} &= S_{ab} = \rho g_{ab} - \frac{1}{2}(\nabla_a \xi_b + \nabla_b \xi_a) = \tilde{\rho} \tilde{g}_{ab} - \frac{1}{2}(\tilde{\nabla}_a \tilde{\xi}_b + \tilde{\nabla}_b \tilde{\xi}_a), \\ \tilde{S}_{au} &= 0 = -\frac{1}{2}(\tilde{\nabla}_a \tilde{\xi}_u + \tilde{\nabla}_u \tilde{\xi}_a), \\ \tilde{S}_{uv} &= 0 = \tilde{\rho} \delta_{uv} - \frac{1}{2}(\tilde{\nabla}_u \tilde{\xi}_v + \tilde{\nabla}_v \tilde{\xi}_u). \end{aligned}$$

Hence we can state, if B is a Ricci soliton, then $M = R^2 \times B$ is a Ricci soliton.

Conversely, suppose that $M = R^2 \times B$ is a Ricci soliton. Then, we get

$$(10) \quad \begin{aligned} S_{ab} = \tilde{S}_{ab} &= \tilde{\rho} g_{ab} - \frac{1}{2}(\partial_a \tilde{\xi}_b + \partial_b \tilde{\xi}_a - 2 \left\{ \begin{matrix} c \\ ab \end{matrix} \right\} \tilde{\xi}_c), \\ \partial_a \tilde{\xi}_u + \partial_u \tilde{\xi}_a &= 0, \\ \tilde{\rho} \delta_{uv} - \frac{1}{2}(\partial_u \tilde{\xi}_v + \partial_v \tilde{\xi}_u) &= 0. \end{aligned}$$

for some constant $\tilde{\rho}$ on M , where $\tilde{\xi}_i$ is a component of the smooth vector field \tilde{V} on M and the range of indices i, j, k, \dots is $\{1, 2, 3, \dots, m + 2\}$. Considering the case $u = v$ in the third equation of (10), we obtain

$$(11) \quad \tilde{\xi}_u = \tilde{\rho}x_u + h_{(u)}^u,$$

where $h_{(u)}^u$ is a function on M which corresponds to $\tilde{\xi}_u$ and the symbol (u) means having no x_u -variable.

From the third equation of (10) and (11), we obtain

$$(12) \quad \partial_u h_{(v)}^v = -\partial_v h_{(u)}^u,$$

from which we see that $\partial_u h_{(v)}^v$ is a function on B . Thus, we can denote

$$(13) \quad \partial_u h_{(v)}^v = H_{(u,v)},$$

where $H_{(u,v)}$ means a function H having no x_u -variable and x_v -variable.

By use of the equations (11) and (13), we obtain

$$(14) \quad h_{(u)}^u = -H_{(u,v)}x_v + K_{(1,2)}^u,$$

and

$$(15) \quad \tilde{\xi}_u = \tilde{\rho}x_u - H_{(u,v)}x_v + K_{(1,2)}^u, (u \neq v),$$

from the equations (11) and (14).

The second equation of (10) and (15) give rise to

$$(16) \quad \tilde{\xi}_a = -\partial_a K_{(1,2)}^1 x_1 - \partial_a K_{(1,2)}^2 x_2 + L_{(1,2)}^a,$$

and the first equation of (10) gives

$$S_{ab} - \tilde{\rho}g_{ab} = -\frac{1}{2}(\partial_a \tilde{\xi}_b + \partial_b \tilde{\xi}_a - 2 \left\{ \begin{matrix} c \\ ab \end{matrix} \right\} \tilde{\xi}_c).$$

Moreover, we get

$$\tilde{V} = (\tilde{\rho}x_1 - H_{(1,2)}x_2 + K_{(1,2)}^1, \tilde{\rho}x_2 - H_{(2,1)}x_1 + K_{(1,2)}^2, -\partial_3 K_{(1,2)}^1 x_1 - \partial_3 K_{(1,2)}^2 x_2 + L_{(1,2)}^3, \dots, -\partial_{m+2} K_{(1,2)}^1 x_1 - \partial_{m+2} K_{(1,2)}^2 x_2 + L_{(1,2)}^{m+2})$$

by means of the equations (15) and (16).

If we take $V = (\xi_3, \dots, \xi_{m+2})$ such that $\xi_a = L_{(1,2)}^a$, then we get

$$S_{ab} - \rho g_{ab} = -\frac{1}{2}(\nabla_a L_{(1,2)}^b + \nabla_b L_{(1,2)}^a),$$

for $\rho = \tilde{\rho}$. Hence we see that if $M = R^2 \times B$ is a Ricci soliton, then B is a Ricci soliton. Thus we get

THEOREM 2.2. *$M = R^2 \times B$ is a Ricci soliton if and only if B is a Ricci soliton.*

In general, if we assume that the Riemannian products $M = B \times F$ is a Ricci soliton, then there exists vector field $X = (\tilde{\xi}_1, \dots, \tilde{\xi}_n, \tilde{\xi}_{n+1}, \dots, \tilde{\xi}_{n+p})$ on M and constant $\tilde{\rho}$ such that $\tilde{S}_{ij} = \tilde{\rho}\tilde{g}_{ij} - (\tilde{\nabla}_i\tilde{\xi}_j + \tilde{\nabla}_j\tilde{\xi}_i)/2$, that is

$$S_{ab} = \tilde{\rho}g_{ab} - (\tilde{\nabla}_a\tilde{\xi}_b + \tilde{\nabla}_b\tilde{\xi}_a)/2,$$

$$\tilde{\nabla}_a\tilde{\xi}_x + \tilde{\nabla}_x\tilde{\xi}_a = 0,$$

$$\bar{S}_{xy} = \tilde{\rho}g_{xy} - (\tilde{\nabla}_x\tilde{\xi}_y + \tilde{\nabla}_y\tilde{\xi}_x)/2.$$

If we take $\rho = \tilde{\rho}$ and $X_B = (\tilde{\xi}_1, \dots, \tilde{\xi}_n)$, then we see that B is a Ricci soliton. Similarly we find F is Ricci soliton with $\bar{\rho} = \tilde{\rho}$ and $X_F = (\tilde{\xi}_{n+1}, \dots, \tilde{\xi}_{n+p})$.

Conversely, if we assume that B is a Ricci soliton with $X_B = (\xi_1, \dots, \xi_n)$ and constant ρ on B , and F is a Ricci soliton with $X_F = (\xi_{n+1}, \dots, \xi_{n+p})$ and constant $\bar{\rho}$ on F . then we can express

$$S_{ab} = \rho g_{ab} - (\nabla_a \xi_b + \nabla_b \xi_a)/2, \quad \bar{S}_{xy} = \bar{\rho} g_{xy} - (\bar{\nabla}_x \xi_y + \bar{\nabla}_y \xi_x)/2.$$

Now we take $\tilde{\rho} = \rho = \bar{\rho}$ and $X = (X_B, X_F)$ on M , then we have

$$\tilde{S}_{ab} = S_{ab} = \tilde{\rho}\tilde{g}_{ab} - (\tilde{\nabla}_a\tilde{\xi}_b + \tilde{\nabla}_b\tilde{\xi}_a)/2,$$

$$\tilde{S}_{ax} = 0 = \tilde{\rho}\tilde{g}_{ax} - (\tilde{\nabla}_a\tilde{\xi}_x + \tilde{\nabla}_x\tilde{\xi}_a)/2,$$

$$\tilde{S}_{xy} = \bar{S}_{xy} = \tilde{\rho}\tilde{g}_{xy} - (\tilde{\nabla}_x\tilde{\xi}_y + \tilde{\nabla}_y\tilde{\xi}_x)/2,$$

that is, $\tilde{S}_{ij} = \tilde{\rho}\tilde{g}_{ij} - (\tilde{\nabla}_i\tilde{\xi}_j + \tilde{\nabla}_j\tilde{\xi}_i)/2$. Hence we see that $M = B \times F$ is a Ricci soliton. Thus we have

THEOREM 2.3. *Let $M = B \times F$ be a Riemannian product of B and F . Then M is a Ricci soliton if and only if B and F are Ricci solitons with $\rho = \bar{\rho}$.*

3. Gradient Ricci solitons in the warped product spaces

Let (B, g) and (F, \bar{g}) be n and p -dimensional Riemannian manifolds with Riemannian metric g and \bar{g} respectively. Then the warped product space $M = B \times_f F$ with a warping function f has the Riemannian

metric $G = \begin{pmatrix} g & 0 \\ 0 & f^2\bar{g} \end{pmatrix}$. Hence the components of Ricci tensors are given by [1, 10, 11]

$$\tilde{S}_{ab} = S_{ab} - \frac{p}{f}\nabla_a f_b,$$

$$\tilde{S}_{ax} = 0,$$

$$\tilde{S}_{xy} = \bar{S}_{xy} - f(\Delta f)\bar{g}_{xy} - (p-1)\|f_e\|^2\bar{g}_{xy},$$

where \tilde{S} , S and \bar{S} are the Ricci tensors of M , B and F respectively, and Δf is the Laplacian of f for g .

Suppose that $M = B \times_f F$ is a gradient Ricci solution. Then, we have

$$\tilde{S}_{ab} = \tilde{\rho}\tilde{g}_{ab} - \tilde{\nabla}_a\tilde{\nabla}_b h = \tilde{\rho}g_{ab} - \nabla_a\nabla_b h,$$

$$\tilde{S}_{ax} = \tilde{\rho}\tilde{g}_{ax} - \tilde{\nabla}_a\tilde{\nabla}_x h = -\partial_a h_x + \frac{f_a}{f}h_x,$$

$$\tilde{S}_{xy} = \tilde{\rho}\tilde{g}_{xy} - \tilde{\nabla}_x\tilde{\nabla}_y h = \tilde{\rho}f^2\bar{g}_{xy} - (\bar{\nabla}_x\bar{\nabla}_y h + f f^c h_c \bar{g}_{xy}),$$

for some constant $\tilde{\rho}$ and some function h on M .

Then we have

$$(17) \quad S_{ab} - \frac{p}{f}\nabla_a f_b = \tilde{\rho}g_{ab} - \nabla_a h_b,$$

$$(18) \quad \partial_a h_x = \frac{f_a}{f}h_x,$$

$$(19) \quad \bar{S}_{xy} = (\tilde{\rho}f^2 + f(\Delta f) + (p-1)\|f_e\|^2 - f f^c h_c)\bar{g}_{xy} - \bar{\nabla}_x\bar{\nabla}_y h.$$

Assume that the partial derivative of h with respect to all x vanishes, then h becomes a function on B . We see that F is an Einstein space because the coefficient of \bar{g}_{xy} in (19) is constant along F . Thus we have

THEOREM 3.1. *Let $M = B \times_f F$ be a gradient Ricci soliton with a potential function h . If the partial derivative of h with respect to all x vanishes, then F becomes an Einstein space.*

If $h_x \neq 0$ for all x , then equation (18) becomes $\frac{\partial_a h_x}{h_x} = \frac{f_a}{f}$. Hence $\partial_a(\ln h_x - \ln f) = 0$, that is we can $\ln \frac{h_x}{f} = l$ for some function l on F . Therefore $h_x = f e^l$, which induces $h = f k$ for some function k on F . Since $h_x \neq 0$ for all x and f depends on B , $k_x \neq 0$. Hence we have $h_a = f_a k$, $\nabla_b h_a = k \nabla_b f_a$ and the equation (17) becomes

$$(20) \quad S_{ab} - \frac{p}{f} \nabla_a f_b - \tilde{\rho} g_{ab} = k \nabla_b f_a.$$

If we assume that $\nabla_b f_a \neq 0$ for some a and b , then it leads to a contradiction in the equation (20) because the left hand side of (20) only depends on B and k the right hand side depends on B and F . Hence we see that $\nabla_b f_a = 0$ for all a and b and that B becomes an Einstein space from the equation (20). Moreover $\nabla_b f_a = 0$ implies Laplacian of f equal to zero. So if B is compact, then f becomes a constant, that is, M is the Riemannian product space. Thus we have

THEOREM 3.2. *Let $M = B \times_f F$ be a gradient Ricci soliton with a potential function h . If $h_x \neq 0$ for all x , then B becomes an Einstein space. In this case $\nabla_b f_a = 0$ for all a and b , and M is the Riemannian product space if B is compact.*

On the other hand, if the partial derivative $h_a = 0$ for all a , then h only depends on F and $f_a h_x = 0$ from (18). Hence f is a constant function or h is a constant function. The fact f is a constant means that M is the Riemannian product of B and F , moreover B is Einstein and F becomes a gradient Ricci soliton due to (17) and (19). If h is a constant function, then we see that the Ricci curvature on B has the form $S_{ab} = \tilde{\rho} g_{ab} + \frac{p}{f} \nabla_a f_b$ and F is Einstein from (19). Moreover if f and h are constants, then the Ricci curvatures of B and F are given by $S_{ab} = \tilde{\rho} g_{ab}$ and $\tilde{S}_{xy} = \tilde{\rho} f^2 \bar{g}_{xy}$, respectively. Since the metric G on M is given by $G = \begin{pmatrix} g & 0 \\ 0 & f^2 \bar{g} \end{pmatrix}$, we see that $\tilde{S} = \tilde{\rho} G$, that is, M is Einstein. Thus we have

THEOREM 3.3. *Let $M = B \times_f F$ be a gradient Ricci soliton with a potential function h . If the partial derivative $h_a = 0$ for all a , then one of the following three cases occurs.*

- (a) M is the Riemannian product of an Einstein space and a gradient Ricci soliton space.
- (b) F is Einstein.
- (c) M is the Riemannian product of two Einstein spaces, and moreover M is an Einstein space.

4. Gradient Yamabe solitons in the Riemannian product manifolds

At first, we consider the gradient Yamabe soliton in the product space $M = R \times B$ with a Riemannian metric \tilde{g} and a potential function h , where (B, g) is an m -dimensional Riemannian manifold. Then we see that

$$(21) \quad \begin{aligned} (\tilde{r} - \tilde{\rho}) &= \partial_1 h_1, \\ \partial_1 h_a &= 0, \\ (\tilde{r} - \tilde{\rho})g_{ab} &= \nabla_a \nabla_b h, \\ \tilde{r} &= r, \end{aligned}$$

where \tilde{r} and r are scalar curvatures of M and B respectively, and the range of indices a, b, c, \dots is $\{2, 3, \dots, m + 1\}$. From the second equation of (21), we see that the potential function h is decomposed into $h(t, x_1, x_2, \dots, x_m) = l(t) + k(x_1, \dots, x_m)$ for functions l on R and k on B . Then the third equation of (21) can be rewritten as $(\tilde{r} - \tilde{\rho})g_{ab} = \nabla_a \nabla_b k$, that is, B is a gradient Yamabe soliton. Moreover, from (21), we see that $\partial_1 l_1 = \partial_1 h_1 = \tilde{r} - \tilde{\rho} = \frac{1}{m} g^{ab} \nabla_a \nabla_b h$. Since $\partial_1 l_1$ only depends on R and the right hand side is a quantity of B , $\tilde{r} - \tilde{\rho}$ becomes constant, that is, $r = \tilde{r}$ is constant.

Conversely, let (B, g) be a gradient Yamabe soliton with a constant scalar curvature r . Then $(r - \rho)g_{ab} = \nabla_a \nabla_b k$ with a potential function k and $\tilde{r} = r$ is constant. Take the function h on M as $h(t, x_1, x_2, \dots, x_m) = (r - \rho)\frac{t^2}{2} + k(x_1, \dots, x_m)$ and $\tilde{\rho} = \rho$. Then we get $(\tilde{r} - \tilde{\rho}) = \partial_1 h_1, \partial_1 h_a = 0,$
 $(\tilde{r} - \tilde{\rho})\tilde{g}_{ab} = \tilde{\nabla}_a \tilde{\nabla}_b h,$
 that is, $M = R^1 \times B$ becomes a gradient Yamabe soliton. Thus we have

THEOREM 4.1. $M = R^1 \times B$ is a gradient Yamabe soliton if and only if B is a gradient Yamabe soliton with a constant scalar curvature.

References

[1] A.Besse, *Einstein manifolds*, Springer-Verlag, Berlin, 1987.
 [2] H. D. Cao, *Geometry of Ricci solitons*, Lecture note, Lehigh Univ., 2008.
 [3] H. D. Cao, *Recent progress on Ricci solitons*, arXiv:0908.2006v1, 2008.
 [4] T. Chave, G. Valent, *On a class of compact and non-compact quasi-Einstein metrics and their renormalizability properties*, Nuclear Phys. B478 (1996), 758–778.

- [5] M. Eminenti, G. La Nave and C. Mantegazza, *Ricci solitons : The equation point of view*, Manuscripta Math. **127** (2008) 345–367.
- [6] M. Fernández-López and E. García-Río, *A remark on compact Ricci solitons*, Math. Ann. **340** (2008) 893–896.
- [7] R. S. Hamilton, *The Ricci flow on surfaces*, Mathematics and general relativity (Santa Cruz CA, 1986), 237-262, Contemp. Math. **71**, American Math. Soc., 1988.
- [8] T. Ivey, *Ricci solitons on compact 3-manifolds*, Differential Geom. Appl. **3** (1993) 301-307.
- [9] T. Ivey, *New examples of complete Ricci solitons*, Proc. AMS **122** (1994) 241–245.
- [10] B. H. Kim, *Warped products with critical Riemannian metric*, Proc. Japan Academy **71** (1995) 117–118.
- [11] B. H. Kim, S.D.Lee, J.H.Choi and Y.O.Lee, *On warped product spaces with a certain Ricci condition*, B. Korean Math. Soc. **50** (2013) (5), 1693–1710.
- [12] L. Ma and V. Miquel, *Remarks on scalar curvature of Yamabe solitons*, Ann. Glob. Ana., Geom. **42** (2012), 195–205.
- [13] G. Perelman, *Ricci flow with surgery on three manifolds*, arXiv:math.DG/0303109.

Sang Deok Lee
Department of Mathematics
Dankook University
Cheonan 31116, Republic of Korea
E-mail: sdlee@dankook.ac.kr

Byung Hak Kim
Department of Applied Mathematics
Kyung Hee University
Yongin 17104, Republic of Korea
E-mail: bhkim@khu.ac.kr

Jin Hyuk Choi
Humanitas college
Kyung Hee University
Yongin 17104, Republic of Korea
E-mail: jinhchoi@khu.ac.kr